



# Intraspecies Signaling between Common Variants of *Pseudomonas aeruginosa* Increases Production of Quorum-Sensing-Controlled Virulence Factors

Dallas L. Mould,<sup>a</sup> Nico J. Botelho,<sup>a</sup>  Deborah A. Hogan<sup>a</sup>

<sup>a</sup>Geisel School of Medicine at Dartmouth, Hanover, New Hampshire, USA

**ABSTRACT** The opportunistic pathogen *Pseudomonas aeruginosa* damages hosts through the production of diverse secreted products, many of which are regulated by quorum sensing (QS). The *lasR* gene, which encodes a central QS regulator, is frequently mutated in clinical isolates from chronic infections, and loss of LasR function (LasR<sup>−</sup>) generally impairs the activity of downstream QS regulators RhIR and PqsR. We found that in cocultures containing LasR<sup>+</sup> and LasR<sup>−</sup> strains, LasR<sup>−</sup> strains hyperproduce the RhIR/RhII-regulated antagonistic factors pyocyanin and rhamnolipids in diverse models and media and in different strain backgrounds. Diffusible QS auto-inducers produced by the wild type were not required for this effect. Using transcriptomics, genetics, and biochemical approaches, we uncovered a reciprocal interaction between wild-type and *lasR* mutant pairs wherein the iron-scavenging siderophore pyochelin produced by the *lasR* mutant induced citrate release and cross-feeding from the wild type. Citrate, a metabolite often secreted in low iron environments, stimulated RhIR signaling and RhII levels in LasR<sup>−</sup> but not in LasR<sup>+</sup> strains. These studies reveal the potential for complex interactions between recently diverged, genetically distinct isolates within populations from single chronic infections.

**IMPORTANCE** Coculture interactions between *lasR* loss-of-function and LasR<sup>+</sup> *Pseudomonas aeruginosa* strains may explain the worse outcomes associated with the presence of LasR<sup>−</sup> strains. More broadly, this report illustrates how interactions within a genotypically diverse population, similar to those that frequently develop in natural settings, can promote unpredictably high virulence factor production.

**KEYWORDS** *Pseudomonas aeruginosa*, RhIR, citrate, intraspecies interactions, *lasR*, pyochelin, quorum sensing

In chronic infections and healthy microbiomes, genetic diversity frequently arises and persists within clonally derived microbial populations, and recent data highlight that heterogeneity within a population can pose challenges to clearance and treatment (1–4). Genotypic and phenotypic complexity is particularly well documented in the chronic lung infections associated with the genetic disease cystic fibrosis (CF), and studies have convincingly demonstrated that within a species, a common set of genes is under selection across strains and hosts (5–11).

Loss-of-function mutations in *Pseudomonas aeruginosa lasR* (LasR<sup>−</sup>) are commonly found in CF isolates and strains from acute infections and from environmental sources (12–16). LasR participates in the regulation of quorum sensing (QS) in conjunction with other transcription factors, including RhIR and PqsR (Mvfr). These regulators have one or more autoinducer ligands: 3-oxo-C<sub>12</sub>-homoserine lactone (3OC12HSL) for LasR, C<sub>4</sub>-homoserine lactone (C4HSL) for RhIR, and hydroxy-alkyl-quinolones (*pseudomonas* quinolone signal [PQS] and hydroxy-heptyl quinolone [HHQ]) for PqsR (17). In the

**Citation** Mould DL, Botelho NJ, Hogan DA. 2020. Intraspecies signaling between common variants of *Pseudomonas aeruginosa* increases production of quorum-sensing-controlled virulence factors. *mBio* 11:e01865-20. <https://doi.org/10.1128/mBio.01865-20>.

**Editor** Joanna B. Goldberg, Emory University School of Medicine

**Copyright** © 2020 Mould et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Deborah A. Hogan, [dhogan@dartmouth.edu](mailto:dhogan@dartmouth.edu).

This article is a direct contribution from Deborah A. Hogan, a Fellow of the American Academy of Microbiology, who arranged for and secured reviews by Amanda G. Oglesby-Sherrouse, University of Maryland School of Pharmacy, and Ajai A. Dandekar, University of Washington.

**Received** 10 July 2020

**Accepted** 17 July 2020

**Published** 25 August 2020

regulatory networks described in widely used *P. aeruginosa* model strains, LasR is an upstream regulator of RhIR and PqsR signaling, and together these regulators control the expression of a suite of genes associated with virulence, including redox-active small-molecule phenazines (18–20), cyanide (21), and rhamnolipid surfactants important for surface motility, biofilm dispersal, and host cell damage (22–24).

Although LasR positively regulates virulence factors, and *lasR* loss-of-function mutants have reduced virulence in infection models, LasR<sup>−</sup> strain culture positivity is correlated with worse disease outcomes in acute and chronic infections (12, 13). There are several possible explanations for this apparent contradiction. LasR<sup>−</sup> clinical isolates (CIs) are frequently found among strains with functional LasR (LasR<sup>+</sup>) where exoproducts can be shared or signal cross-feeding can occur (14), and some LasR<sup>−</sup> clinical isolates exhibit rewired QS regulation (25). Loss of LasR function also confers some fitness advantages, including altered catabolic profiles (26) and enhanced growth in low oxygen (27, 28), which may contribute to bacterial burden. Further, LasR<sup>−</sup> strains can activate QS in response to specific fungal products (29) or culture conditions (30, 31).


In addition to LasR status, iron acquisition strategies are often heterogeneous across *P. aeruginosa* isolates. *P. aeruginosa* procures iron through the use of siderophores, including pyochelin (32–34) and pyoverdine (35), from heme, or through a direct iron uptake system (36–38). Although it is common to encounter *P. aeruginosa* strains with loss-of-function mutations in genes required for biosynthesis of the high-affinity siderophore pyoverdine, genes associated with use of pyochelin, heme utilization, and ferrous iron import are generally intact (39–41). Iron limitation can deprioritize pathways that require abundant iron including the tricarboxylic acid (TCA) cycle (42), and consequently *Pseudomonas* spp. and other organisms release metabolic intermediates, such as citrate, that accumulate at iron-requiring steps (e.g., aconitase) (43, 44).

Here, we show that mixtures of *P. aeruginosa* LasR<sup>−</sup> and LasR<sup>+</sup> strains had enhanced production of QS-controlled factors across media, culture conditions, and strain backgrounds. The unpredictably high levels of exoproducts in coculture were produced by LasR<sup>−</sup> strains due to activation of RhIR, likely through increased C4HSL synthase (RhII) stability in LasR<sup>−</sup> strains. Our genetic, transcriptomic, and biochemical studies led us to uncover a set of interactions in which production of the siderophore pyochelin by  $\Delta$ *lasR* cells induced citrate release by wild type (WT) but not by  $\Delta$ *lasR* cells. We found that citrate led to increased RhII protein levels and RhIR activity in  $\Delta$ *lasR* cells but not in the WT. Together, these intraspecies interactions increased production of exoproducts known to cause host damage.

## RESULTS

### *P. aeruginosa* $\Delta$ *lasR* overproduces pyocyanin in coculture with the wild type.

We observed that mixtures of *P. aeruginosa* LasR<sup>+</sup> and LasR<sup>−</sup> strains had high levels of total pyocyanin, a secreted, blue-pigmented phenazine. As shown in spot colonies of the PA14 wild-type (WT),  $\Delta$ *lasR* strain, and WT and  $\Delta$ *lasR* strain cocultures (here, WT/ $\Delta$ *lasR* cocultures), the strain mixture showed increased blue pigmentation (Fig. 1A) and a significant 4-fold induction of pyocyanin above the background relative to the level with either strain alone (Fig. 1B). Phenazine-deficient derivatives,  $\Delta$ *phz* ( $\Delta$ *phzA1*  $\Delta$ *phzB1*  $\Delta$ *phzC1*  $\Delta$ *phzD1*  $\Delta$ *phzE1*  $\Delta$ *phzF1*  $\Delta$ *phzG1*  $\Delta$ *phzA2*  $\Delta$ *phzB2*  $\Delta$ *phzC2*  $\Delta$ *phzD2*  $\Delta$ *phzE2*  $\Delta$ *phzF2*  $\Delta$ *phzG2*) and  $\Delta$ *lasR*  $\Delta$ *phz* ( $\Delta$ *lasR*  $\Delta$ *phzC1*  $\Delta$ *phzC2*) strains, were also included, and as expected,  $\Delta$ *phz* and  $\Delta$ *lasR*  $\Delta$ *phz* cells showed no blue colony pigmentation (Fig. 1A). The higher levels of pyocyanin in WT/ $\Delta$ *lasR* cocultures relative to levels in single-strain cultures were also observed on artificial sputum medium (ASM) and on phosphate-buffered medium with or without amino acids, indicating that the phenomenon occurred under diverse conditions (see Fig. S1A in the supplemental material). Cocultures of clonally derived LasR<sup>+</sup> and LasR<sup>−</sup> clinical isolates collected from single respiratory sputum samples from chronically infected individuals with cystic fibrosis (14) also had increased production of pyocyanin relative to monoculture levels when LasR<sup>+</sup> and LasR<sup>−</sup> strains were grown together (Fig. S1B and C).



**FIG 1** The  $\Delta lasR$  strain produces pyocyanin in wild-type/ $\Delta lasR$  cocultures beyond monoculture concentrations. (A) Representative images of the wild-type (WT) and  $\Delta lasR$  strains and their phenazine-deficient derivatives ( $\Delta phz$  strains) visualized from the bottom of a 96-well LB agar plate after 16 h growth as mono- and cocultures with 70:30 WT-to- $\Delta lasR$  cell initial ratio. (B) Pyocyanin levels above background, defined as the average signal for the  $\Delta lasR$   $\Delta phz$  strain, quantified for cultures described in panel A. ns, not significant; \*,  $P < 0.05$ ; \*\*\*,  $P < 0.0005$ ; \*\*\*\*,  $P < 0.0001$ , as determined by ordinary one-way analysis of variance with Tukey's multiple-comparison test for  $n \geq 8$  replicates on three different days. (C) Model of pyocyanin production by the  $\Delta lasR$  strain in coculture with the WT. (D) Representative pyocyanin production of the wild type cocultured with the  $\Delta lasR$  strain or the  $\Delta lasR$  strain complemented with the  $lasR$  gene at the native locus ( $\Delta lasR$  +  $lasR$  strain) across several initial (designated by the subscript i) proportions on LB medium for 20 h. Three biological replicates were included in at least 3 independent experiments. (E) Average final proportion of 3 replicate colony biofilms quantified after 16 h growth for the WT strain and the  $\Delta lasR$  strain cocultured with a WT strain tagged with  $lacZ$  in 3 independent experiments. Experimental setup was as described in panel D. \*,  $P < 0.05$ ; \*\*,  $P < 0.005$ , as determined by two-tailed t tests of paired ratios between  $att::lacZ$ /WT (control) and  $att::lacZ$ / $\Delta lasR$  cocultures at each initial ratio. All results that reach significance are marked.

To assess individual strain contributions to increased pyocyanin in WT/ $\Delta lasR$  cocultures, we replaced each strain with its phenazine-deficient derivative and measured pyocyanin in coculture. When the  $\Delta lasR$  strain was cultured with the phenazine biosynthesis mutant  $\Delta phz$  strain ( $\Delta phz$ / $\Delta lasR$  coculture), we still observed increased blue pigmentation and total pyocyanin at a level above that of either monoculture (Fig. 1A and B). Surprisingly, pyocyanin production by  $\Delta phz$ / $\Delta lasR$  cocultures was statistically higher than that of WT/ $\Delta lasR$  coculture (Fig. 1B). In contrast, WT/ $\Delta lasR$   $\Delta phz$  cocultures did not display the high-pyocyanin phenotype (Fig. 1A and B) and resembled  $\Delta phz$ / $\Delta lasR$   $\Delta phz$  cocultures, where no pyocyanin was produced. Collectively, these data suggested that WT/ $\Delta lasR$  cocultures produced more pyocyanin than either monoculture alone and that the  $\Delta lasR$  strain contributed the pyocyanin in coculture (Fig. 1C).

That the levels of pyocyanin in WT/ $\Delta lasR$  cocultures were higher than the level in each strain alone was not dependent on the initial ratios of WT to  $\Delta lasR$  cells (Fig. 1D). We saw increased coculture colony pigmentation when the initial proportions of WT

cells were at 0.2, 0.3, 0.5, 0.7, and 0.8 of the initial inoculums, with the balance comprised of  $\Delta lasR$  cells (Fig. 1D).


No increase in pyocyanin was observed at any ratio when the WT was cocultured with the  $\Delta lasR$  complemented derivative strain ( $\Delta lasR + lasR$ ), indicating that the phenomenon was dependent on the *lasR* mutation (Fig. 1D). To assess the relative abundances of WT and  $\Delta lasR$  cells in coculture, we competed each strain against a neutrally tagged WT strain (PA14 *att::lacZ*). We found that  $\Delta lasR$  cells increased in proportion after 16 h of growth in colony biofilms regardless of the starting proportion whereas the proportions of untagged WT cells remained stable (Fig. 1E). We have previously shown that Anr activity is higher in  $\Delta lasR$  strains and contributes to the competitive fitness of the  $\Delta lasR$  strain against WT *P. aeruginosa* in colony biofilms (27, 45), but Anr was not required for coculture pyocyanin production (Fig. S2).

Pyocyanin is a product regulated by quorum sensing (QS) through the transcription factors LasR, RhIR, and PqsR (46–48), and because QS regulation is cell density dependent, it was important to assess the coculture population size relative to that of the monoculture. Total CFU counts did not increase in WT/ $\Delta lasR$  mixed cultures relative to the level for either strain alone (Fig. S1D). Instead, we found that WT/ $\Delta lasR$  cocultures had fewer CFU than WT monocultures on lysogeny broth (LB) medium (Fig. S1D). Taken together, these data suggested that altered behavior, rather than cell number, contributed to the increased phenazine profile of LasR<sup>−</sup> strains.

**Independent of its ability to produce autoinducers, the WT promotes RhIR/I-dependent signaling in a  $\Delta lasR$  strain.** In the canonical QS pathway, LasR regulates both PqsR and RhIR, and mutants lacking either regulator in a WT background have impaired pyocyanin production (49, 50). Both *pqsR* and *rhIR* were required in  $\Delta lasR$  cells for pyocyanin production in coculture with the WT (Fig. 2A). To determine if coculture increased RhIR- or PqsR-dependent signaling in  $\Delta lasR$  strains, we fused *lacZ* to the promoters of *rhII* and *pqsA* (*PrhII* and *PpqsA*, respectively) which provide activity readouts of each respective regulator (17). We examined the interactions between WT and the  $\Delta lasR$  strain in single-cell-derived colonies by spreading suspensions containing ~50 cells of WT with ~50 cells of either a  $\Delta lasR$  *PrhII-lacZ* or  $\Delta lasR$  *PpqsA-lacZ* strain on LB agar containing the colorimetric  $\beta$ -galactosidase substrate 5-bromo-4-chloro-3-indolyl-D-galactopyranoside (X-Gal). Intercolony distances and  $\beta$ -galactosidase activity in  $\Delta lasR$  strains were measured. We found that the rise in *PrhII-lacZ* activity was inversely correlated with the distance to a WT colony (Fig. 2B). Pearson correlation analyses showed that 54% of the variability in  $\Delta lasR$  *PrhII-lacZ* strain activity could be explained by changes in the distance to a WT colony (*P* value of  $\leq 0.0001$ ). The increased *PrhII-lacZ* activity in the  $\Delta lasR$  strain was not observed in the  $\Delta lasR \Delta rhIR$  strain, and close proximity to another  $\Delta lasR$  *PrhII-lacZ* colony did not affect promoter activity (Fig. 2B, inset). Because C4HSL (which is synthesized by RhII) activates RhIR and because proximity to the WT stimulated  $\Delta lasR$  *PrhII-lacZ* strain activity, we examined the role of RhII in the  $\Delta lasR$  strain response. We observed that a  $\Delta lasR \Delta rhII$  strain was greatly impaired in the induction of pyocyanin upon coculture with the WT (Fig. 2A), which suggests that WT production of C4HSL was insufficient to complement the  $\Delta lasR \Delta rhII$  strain and further posits activation of RhIR and C4HSL synthesis in  $\Delta lasR$  strains. Although PqsR was required in  $\Delta lasR$  cells for coculture pyocyanin production, there was no significant correlation with proximity to the WT for  $\Delta lasR$  *PpqsA-lacZ* strain activity (Fig. 2B). Collectively, these data indicated that a diffusible factor produced by the WT stimulated RhIR-dependent signaling in the  $\Delta lasR$  strain to induce downstream production of RhIR- and PqsR-dependent factors.

Given differences in colony pigmentation between WT/ $\Delta lasR \Delta rhIR$  and WT/ $\Delta lasR \Delta rhII$  (Fig. 2A) cocultures, C4HSL cross-feeding between the WT and  $\Delta lasR$  strain likely occurred. Because C4HSL is diffusible and produced by WT cells, we tested the hypothesis that C4HSL or other acyl-homoserine lactones (AHLs) produced by the WT were necessary to induce RhIR-dependent activity in  $\Delta lasR$  cells cocultured with the WT. To test this hypothesis, we cocultured the  $\Delta lasR$  strain with  $\Delta rhII$  cells or  $\Delta lasI \Delta rhII$  cells which lack both acyl-homoserine lactone synthases. Surprisingly, we found that like





**FIG 2** *P. aeruginosa* WT induces RhIR/I-dependent pyocyanin production in  $\Delta lasR$  cells even in the absence of WT autoinducers. (A) Representative pyocyanin production by monocultures and WT cocultures (70% WT at time 0) of  $\Delta lasR$  and  $\Delta lasR$  strain derivatives that are deficient in PQS or RhIR/I-dependent quorum sensing on LB medium after 24 h growth. (B) Promoter activity of the  $\Delta lasR$   $PpqsA-lacZ$  (gray) and  $\Delta lasR$   $PrhII-lacZ$  (black) strains, quantified by relative pixel intensity of single-cell-derived  $\Delta lasR$  colonies grown near unmodified WT colonies. Solid best-fit nonlinear lines are for visualization. The inset shows representative colonies for RhIR-dependent  $\Delta lasR$   $PrhII-lacZ$  strain activity in monoculture and coculture with the WT (red circles). The experiment was repeated with 2 replicates on at least 3 independent days. (C) Representative monoculture and  $\Delta lasR$  coculture images for the  $\Delta pqsA$ ,  $\Delta rhII$ , and  $\Delta lasI \Delta rhII$  mutants on LB medium after 24 h and 48 h. (30%  $\Delta lasR$  cells at time 0). (D) Model of AHL-dependent and -independent induction of RhIR/I-dependent activity and phenazine production in  $\Delta lasR$  cells grown in coculture with the WT.


WT/ $\Delta lasR$  cocultures,  $\Delta rhII/\Delta lasR$  cocultures had higher levels of pyocyanin than monocultures (Fig. 2C). Similarly,  $\Delta lasI \Delta rhII/\Delta lasR$  cocultures had higher levels of pyocyanin production than monocultures though the interaction was delayed by ~24 h relative to the interaction of the WT/ $\Delta lasR$  cocultures (Fig. 2C). Consistent with the activity of the  $\Delta lasR$   $PpqsA-lacZ$  strain, which was not induced in coculture with WT, the PQS-deficient  $\Delta pqsA$  strain supported high pyocyanin colony pigmentation in coculture with  $\Delta lasR$  cells after 24 h of extended incubation (Fig. 2C). The AHL-independent activation observed in  $\Delta lasI \Delta rhII/\Delta lasR$  cocultures and the striking differences in pyocyanin production observed between the strongly stimulating  $\Delta rhII/\Delta lasR$  cocultures and the weakly stimulating WT/ $\Delta lasR$   $\Delta rhII$  cocultures suggested that the  $\Delta lasR$  strain may rely more heavily on production of its own autoinducer for activation in coculture. Consistent with this model, we found that the  $\Delta lasR$  strain produces RhIR/RhII (RhIR/I)-dependent AHLs in coculture with an AHL-sensing reporter strain (i.e.,  $\Delta lasI \Delta rhII$  strain with a  $lacZ$  promoter fusion to an AHL-responsive gene) (Fig. S3A and B). The dispensable contribution of WT-produced autoinducers implicated a novel signaling interaction in coculture-dependent activation of RhIR/I activity in the  $\Delta lasR$  strain (Fig. 2D).

To assess whether WT-induced RhIR activity in the  $\Delta lasR$  strain was sufficient to elicit other RhIR/I-controlled phenotypes in addition to pyocyanin production, we tested whether coculture with LasR+ strains enhanced swarming, a surface-associated motility which requires the production of RhIR-regulated rhamnolipid surfactants (51). While

the rhamnolipid-defective mutant  $\Delta rhIA$ ,  $\Delta lasR$ , and  $\Delta lasR \Delta rhIR$  strains were unable to swarm, cocultures of the  $\Delta lasR$  strain with the  $\Delta rhIA$  strain swarmed considerably. The phenomenon was dependent on RhlR as the  $\Delta rhIA/\Delta lasR \Delta rhIR$  cocultures did not swarm (Fig. S4). Altogether, these data implicated broad activation of RhlR-mediated QS in LasR<sup>−</sup> strains cocultured with LasR<sup>+</sup> *P. aeruginosa*.

**Pyochelin production by  $\Delta lasR$  cells is required for coculture interactions.** With evidence indicating that induction of RhlR activity in  $\Delta lasR$  cells can occur in both mixed-strain spot colonies and adjacent colonies independent of autoinducer cross-feeding, we sought to gain further insight into the mechanisms that underlie the WT- $\Delta lasR$  cell interactions. We investigated the transcriptomes of the *lasR* mutant in coculture with either the WT or itself via RNA sequencing (RNA-seq). We grew  $\Delta lasR$  colony biofilms on LB medium physically separated from a lawn of either the  $\Delta lasR$  or WT strain by two filters with 0.22- $\mu$ m pores to prevent mixing of genotypes while allowing for the passage of small molecules. In order to examine  $\Delta lasR$  strain transcriptional profiles, RNA was extracted from cells within the  $\Delta lasR$  colony biofilms grown on the topmost filter for 16 h (Fig. 3A). As expected, no *lasR* reads were detected in our sequencing data to suggest that the wild type was sufficiently excluded by filter separation. Expression levels of a total of 199 genes in the  $\Delta lasR$  strain were higher, and those of 198 genes were lower by a  $|\log_2(\text{fold change})|$  of  $\geq 1$  with a *P* value of  $<0.05$ , in coculture with the WT than levels in the  $\Delta lasR$  strain alone (Table S1). Gene Ontology (GO) term analyses through PantherDB (52) indicated that the upregulated gene set was significantly overrepresented in two pathways related to siderophore biosynthesis: the pyoverdine biosynthetic process and salicylic acid biosynthetic process (an upstream precursor of pyochelin) with  $\sim 44$ - and  $\sim 77$ -fold enrichment, respectively (*P* values of  $<0.005$ ). Twenty-eight out of the 33 genes in the pyochelin and pyoverdine siderophore biosynthesis- and acquisition-related GO families were significantly upregulated in  $\Delta lasR$  cells upon coculture with the WT (Fig. 3B) (i.e.,  $|\log_2(\text{fold change})|$  of  $\geq 0$  with a *P* value of  $<0.05$ ). Other low-iron-responsive genes were differentially expressed, including the *has* genes involved in heme uptake and *antABC* genes (Table S1). While we observed stimulation of *rhlI* promoter activity and increased production of RhlR-regulated products, we did not see a broad transcriptional pattern indicative of RhlR activation at this early time point (Table S1), and this point is discussed below.

Given that siderophore biosynthesis genes were upregulated in  $\Delta lasR$  cells cocultured with WT cells, we qualitatively examined production of fluorescent pyoverdine and pyochelin siderophores in monocultures and cocultures. To determine the contribution of both pyoverdine and pyochelin by the  $\Delta lasR$  strain to fluorescence, genes required for pyoverdine biosynthesis ( $\Delta lasR \Delta pvdA$  strain), pyochelin biosynthesis ( $\Delta lasR \Delta pchE$  strain), or both pathways ( $\Delta lasR \Delta pvdA \Delta pchE$  strain) were disrupted (Fig. 3C). Increased fluorescence attributable to both pyoverdine and pyochelin in coculture was due to siderophore production by  $\Delta lasR$  strains, consistent with the RNA-seq data, as the increased fluorescence in WT/ $\Delta lasR$  cocultures was lost in coculture when the  $\Delta lasR$  strain was replaced with a  $\Delta lasR \Delta pvdA$ ,  $\Delta lasR \Delta pchE$ , or  $\Delta lasR \Delta pvdA \Delta pchE$  mutant. While cocultures of the WT and the pyoverdine-deficient derivative  $\Delta lasR \Delta pvdA$  strain (i.e., WT/ $\Delta lasR \Delta pvdA$  coculture) showed increased pyocyanin production relative to that of either monoculture, the  $\Delta lasR \Delta pchE$  and  $\Delta lasR \Delta pvdA \Delta pchE$  strains did not promote pyocyanin production in coculture with the WT, as observed by colony pigmentation (Fig. 3D). The decrease in  $\Delta lasR$  strain-derived pyocyanin was not due to decreased fitness as disruption of *pvdA* and *pchE* individually in  $\Delta lasR$  cells had no effect on the final proportions; in contrast, the  $\Delta lasR \Delta pvdA \Delta pchE$  strain had a significant defect in competitive fitness compared to the fitness of the  $\Delta lasR$  parental strain (Fig. S5). These data suggested that pyochelin played a role in the coculture interaction. To test this model, we complemented the pyocyanin defect in the siderophore-deficient  $\Delta pvdA \Delta pchE/\Delta lasR \Delta pchE$  coculture with pyochelin-containing extracts from cultures of  $\Delta pvdA$  cells which cannot produce pyoverdine or control extracts from



**FIG 3** Biosynthesis of the coculture-induced iron scavenging siderophore pyochelin is required in the  $\Delta lasR$  strain for pyocyanin production when it was cultured with the wild type (WT). (A) Scheme for the collection of RNA from  $\Delta lasR$  colony biofilms grown above a lawn of  $\Delta lasR$  or WT cells. DE, differential expression. (B) Volcano plot showing differential expression ( $\log_2$ ) for  $\Delta lasR$  cells grown over WT relative to  $\Delta lasR$  cells grown over  $\Delta lasR$  on the x axis; the y axis shows the  $-\log_{10} P$  value for the difference between sample types. Genes involved in pyoverdine (blue) and pyochelin (green) iron acquisition systems are indicated. *ccmC* and *ccmF* (indicated with arrows) of the pyoverdine GO term are involved in c-type cytochrome biosynthesis, and strains with knockouts of these genes are reported to produce more pyochelin. (C) Monocultures and WT cocultured with  $\Delta lasR$  strains deficient in pyoverdine ( $\Delta pvdA$ ) and/or pyochelin ( $\Delta pchE$ ) biosynthesis. Colonies were visualized under UV light in order to see fluorescent siderophores. Images are representative of at least 3 independent experiments. (D) Pyocyanin production visualized for the colonies shown in panel C. (E) Representative pyocyanin production by siderophore-deficient strains grown in mono- and coculture on LB medium or on LB medium with pyochelin-containing extract (+PCH). Colonies were grown in a 12-well plate with a 2-ml total volume and imaged after 24 h. (F) Mixed colony biofilms of wild-type and  $\Delta lasR$  strains grown on LB medium (–) or LB medium supplemented with either 10 or 100  $\mu M$   $FeSO_4$  visualized under ambient (top) and UV (bottom) light. (G) Model showing that pyochelin (PCH) production by the  $\Delta lasR$  strain is required for WT/ $\Delta lasR$  coculture phenazines.

siderophore-deficient  $\Delta pvdA \Delta pchE$  cultures (Fig. S6A gives supernatant absorption spectra). The two extracts were analyzed using a chrome azurol S (CAS) assay (53) to confirm that chelator activity was present in the  $\Delta pvdA$  cell supernatant extracts but not in extracts from  $\Delta pvdA \Delta pchE$  cultures (Fig. S6B). Medium supplemented with pyochelin-containing extracts, but not siderophore-free extracts, restored pyocyanin production in  $\Delta pvdA \Delta pchE/\Delta lasR \Delta pchE$  cocultures (Fig. 3E), lending further support to the idea that pyochelin was required for coculture interactions. Consistent with this requirement, iron supplementation suppressed siderophore production, as expected, and diminished coculture pyocyanin in WT/ $\Delta lasR$  cocultures (Fig. 3F) alongside a

decrease in  $\Delta lasR$  strain RhIR/I-dependent AHL activity in coculture with the AHL-sensing reporter strain (Fig. S3C and D). Collectively, these data support a model in which pyochelin production by the  $\Delta lasR$  strain is induced and required for pyocyanin-promoting interactions with the WT through initiation of a low-iron response (Fig. 3G).

**In coculture with the WT, the  $\Delta lasR$  strain responds to citrate, a pyochelin-inducible metabolite.** Many of the upregulated genes in the  $\Delta lasR$  strain upon coculture with the WT have annotations related to organic acids, such as anthranilate and citrate (Table S1 and Fig. S7A). Several lines of evidence suggest that anthranilate was not the factor that stimulated RhIR activity and pyocyanin production in coculture. First, anthranilate supplementation (up to  $\sim 15$  mM) did not alter  $\Delta lasR$  strain phenazine production (Fig. S7B). Further, cocultures of the  $\Delta lasR$  mutant with the anthranilate synthase mutant  $\Delta phnAB$  strain, with reduced extracellular anthranilate (Fig. S7C), or with the  $\Delta pqsA$  mutant (Fig. 2C), which accumulates it (54), did not alter coculture phenazine production. Anthranilate is also generated from tryptophan catabolism through the kynurenine pathway (55). Given that coculture pyocyanin production could occur in the absence of exogenous amino acids (Fig. S1A), we concluded that the kynurenine pathway was likely not involved. Together, these data suggested that anthranilate was not a stimulating metabolite.


In light of the observation that 20% of the most strongly differentially expressed genes [ $|\log_2$  (fold change)| of  $\geq 2$  with a  $P$  value of  $< 0.05$ ] were implicated in citrate sensing, transport, catabolism, and anabolism, as annotated by UniProt (56) and [www.pseudomonas.com](http://www.pseudomonas.com) (57), we looked at all genes with annotations related to citrate to identify broad expression patterns (Fig. 4A). Among the genes induced in  $\Delta lasR$ /WT cocultures [ $|\log_2$  (fold change)| of  $\geq 0$  with a  $P$  value of  $< 0.05$ ] were genes annotated as citrate responsive or playing roles in citrate sensing and transport or metabolism, with the most strongly upregulated citrate genes involved in sensing, transport, and catabolism specifically (Fig. 4B).

We measured citrate in the supernatants of WT and  $\Delta lasR$  LB cultures based on the following observations: (i)  $\Delta lasR$  strains induced low-iron-responsive genes when grown near the WT but not itself; (ii)  $\Delta lasR$  strain pyochelin production was necessary for coculture interactions that led to increased pyocyanin; (iii) citrate sensing and catabolism genes were induced in  $\Delta lasR$  cells by the presence of WT cells; and (iv) numerous microbes, including *Pseudomonas putida*, were shown to secrete citrate and other organic acids when iron limited (44, 58–60). Citrate was detected in both WT and  $\Delta lasR$  strain LB culture supernatants (Fig. 4C), and amendment with extracts containing 50  $\mu$ M pyochelin increased extracellular citrate concentrations by  $\sim 2$ -fold in WT cultures compared to levels in cultures supplemented with extracts lacking pyoverdine and pyochelin, with a much smaller stimulation in  $\Delta lasR$  cultures under the same conditions (Fig. 4C). This suggested that WT-produced citrate may be involved in WT/ $\Delta lasR$  coculture interactions and that citrate release was enhanced by pyochelin produced by the  $\Delta lasR$  strain.

**Citrate induces RhIR-dependent activity and RhII levels in a ClpX protease-dependent manner in  $\Delta lasR$  cells.** To determine if citrate was sufficient to stimulate RhIR activity in  $\Delta lasR$  cells, we analyzed its effects on *rhII* promoter fusion activity, colony morphology, and RhII protein levels. We found that citrate increased *rhII* promoter activity (*PrhII*) in  $\Delta lasR$  cells and that its effects were dependent on the presence of RhIR (Fig. 5A). Citrate was sufficient to promote increases in colony pigmentation and colony smoothness, previously characterized to be RhIR-mediated in  $\Delta lasR$  cells (29) (Fig. 5A, inset). In contrast, citrate caused a small but significant reduction in WT *PrhII* activity compared to that the LB control (Fig. 5A).


To determine if RhII protein levels were influenced by citrate, we utilized an arabinose-inducible *rhII*-hemagglutinin (HA) construct to assess RhII protein levels and stability in the absence and presence of citrate independent of RhIR transcriptional control. RhII-HA was functional as swarming defects of the  $\Delta rhII$  mutant were complemented upon expression of RhII-HA but not by the empty vector (Fig. 5B, inset). RhII-HA protein levels were 3-fold higher in the  $\Delta lasR$  strain upon citrate supplementation than





**FIG 4** Citrate release by WT is induced by pyochelin exposure. (A) Subset of expression data in  $\Delta lasR$  cocultures (Fig. 3A shows the setup) for genes annotated as being involved in citrate sensing, transport, catabolism, anabolism, and those shown to be responsive to citrate. (B) Volcano plot of expression data of  $\Delta lasR$  cocultures with each point representing the  $\log_2(\Delta lasR$  cells grown on WT/ $\Delta lasR$  cells grown on  $\Delta lasR$  cells) expression and  $-\log_{10}(P$  value) of a single gene. Genes shown in panel A are highlighted in color. (C) Citrate concentrations in supernatants from wild-type and  $\Delta lasR$  stationary-phase cultures after growth in LB medium supplemented with extracts containing 50  $\mu\text{M}$  pyochelin (PCH+) or an equal volume of control extracts (PCH-). A representative experiment with four biological replicates repeated 3 independent days is shown. \*\*\*,  $P < 0.0005$ ; \*\*\*\*,  $P \leq 0.0001$ , as determined by two-tailed  $t$  tests of paired ratios.


in the controls (Fig. 5B). Consistent with the absence of an increase in *rhII* promoter activity in WT strains (Fig. 5A), RhII-HA protein levels were not higher with citrate in the  $\Delta lasR$  complemented strain ( $\Delta lasR + lasR$  strain) (Fig. 5B). The differential responses to citrate were also observed in LasR<sup>−</sup> and LasR<sup>+</sup> pairs of clinical isolates (CIs). LasR<sup>−</sup> CIs from acute (strain 388D) and chronic (strain DH2415) infections had RhII-HA levels 1.4- and 1.7-fold higher, respectively, in the presence of citrate (Fig. 5C), whereas alterations in RhII-HA protein levels in LasR<sup>+</sup> CIs from acute (strain 550A) or chronic (strain DH2417) infections were not observed (Fig. 5D). Through this work, we successfully identified citrate as a molecule in coculture that specifically promoted RhII protein levels in LasR<sup>−</sup> strains, but not in LasR<sup>+</sup> strains, by posttranscriptional control. In an attempt to identify transporters involved in the  $\Delta lasR$  strain response to citrate and/or other coculture metabolites, we deleted two organic acid transporters: *dctA* (61) and PA14\_51300 (62) in the  $\Delta lasR$  strain background. We found that the  $\Delta lasR \Delta dctA$  strain showed induction of pyocyanin when it was cocultured with the WT and that induction was dependent on RhIR (Fig. S7D). Similar results were obtained with the  $\Delta lasR$



**FIG 5** Citrate induced RhIR-dependent *rhII* promoter activity and stabilized RhII protein in LasR<sup>-</sup> strains in a ClpX protease-dependent manner. (A)  $\beta$ -Galactosidase activity for  $\Delta lasR$ ,  $\Delta lasR \Delta rhIR$ , and wild-type strains harboring *att::RhII-lacZ* on LB medium with or without 20 mM citrate at 24 h. Each point is the average of three biological replicates from 3 to 4 independent experiments. ns, not significant; \*,  $P < 0.05$ ; \*\*,  $P < 0.005$ , as determined by one-way analysis of variance with Dunnett's multiple hypotheses correction of the indicated comparisons. The inset (A<sub>i</sub>) shows a representative image of  $\Delta lasR$  colony morphology on LB medium with and without 20 mM citrate at 24 h. (B) RhII-HA protein signal normalized to Revert total protein stain (LiCor) on LB medium with and without 20 mM citrate for the  $\Delta lasR$  strain and *lasR* complemented at the native locus ( $\Delta lasR$  strain). ns, not significant; \*,  $P < 0.05$ ; \*\*,  $P < 0.005$ , as determined by analysis of variance with Dunnett's multiple hypothesis correction for  $n = 3$  biological replicates performed on three independent days. The inset (B<sub>i</sub>) illustrates that plasmid-borne RhII-HA, but not the empty vector (EV), complemented an  $\Delta rhII$  mutant for swarming. (C) RhII-HA protein levels on LB medium with and without 20 mM citrate of LasR loss-of-function (LasR<sup>-</sup>) clinical isolates (CIs) from acute corneal (388D) and chronic CF (DH2415) infections. \*,  $P < 0.05$ , as determined by two-tailed *t* tests of paired ratios for  $n = 3$  experiments for each isolate. (D) RhII-HA protein levels on LB medium with and without 20 mM citrate of an LasR<sup>+</sup> acute corneal CI (550A) of the same multilocus sequence type as 388D and LasR<sup>+</sup> chronic CF CI (DH2417) from which DH2415 evolved. ns, not significant as determined by two-tailed *t* tests of paired ratios for  $n = 3$  experiments for each isolate. nd, not detected. (E) Representative image and quantification of replicates for the anti-HA antibody analysis of  $\Delta lasR$  and  $\Delta lasR clpX::TnM$  strains carrying a plasmid expressing RhII-HA or an empty vector (pRhII-HA or pEV, respectively) grown in LB medium with or without 20 mM citrate. \*,  $P < 0.05$ , as determined by two-tailed *t* tests of paired ratios for  $n = 3$  replicates from 3 independent days.

ΔPA14\_51300 strain (Fig. S7E), suggesting that these transporters were not required for the interaction, perhaps due to redundant functions of other proteins or the involvement of other import mechanisms.

The temporal pattern of activation and the stimulation of RhII-HA in the absence of RhIR control suggested that the RhII protein may precede signal amplification via the



**FIG 6** Model for wild-type and  $\Delta lasR$  coculture interactions.  $\Delta lasR$  strain-produced pyochelin promotes citrate release in the wild type (1). Citrate (and diffusible autoinducer) released by the wild type in coculture stimulates RhIR/I-dependent activity in a  $\Delta lasR$  strain (2). Citrate stabilizes RhII protein in  $\Delta lasR$  cells potentially through a ClpXP protease-dependent mechanism (3), which ultimately promotes the production of antagonistic factors like pyocyanin toxin and rhamnolipid surfactant above monoculture levels (4).

QS transcriptional network. This would be consistent with a primary effect on post-transcriptional modulation of RhII-mediated RhIR activity, as has been reported previously to occur through an RhIS small RNA (sRNA)-dependent mechanism (63); however, we found no apparent difference in RhIS expression levels in our RNA-seq reads in coculture. To begin to unravel the mechanisms by which citrate promoted RhIR/I-dependent signaling and RhII stability in  $\Delta lasR$  cells, we analyzed the role of two proteases previously found to target and degrade RhII (i.e., Lon and ClpXP) (64). Given that knockouts of Lon protease have a less substantial rise in RhIR/I expression in  $\Delta lasR$  strains than in the WT (65), we focused on the role of ClpXP in  $\Delta lasR$  cells. We found that citrate induction of RhII-HA protein levels in the  $\Delta lasR$  strain relative to that in the LB control was dependent on functional ClpX protease (Fig. 5E). More specifically, when ClpX, a protease shown to degrade RhII, is nonfunctional (i.e.,  $\Delta lasR$  *clpX::TnM* strain), RhII-HA levels did not increase on LB plus citrate relative to that of the LB control, unlike the level in the  $\Delta lasR$  strain comparator (Fig. 5E). Under LB culture conditions, RhII-HA levels were  $3.20 \pm 2.1$ -fold higher in the  $\Delta lasR$  *clpX::TnM* strain than in the  $\Delta lasR$  strain, which mirrors the 3-fold induction observed for the  $\Delta lasR$  strain on LB medium with or without citrate. Under citrate-supplemented conditions, no significant difference in RhII-HA levels was observed for the  $\Delta lasR$  *clpX::TnM* relative to that in the  $\Delta lasR$  strain (fold change of  $1.01 \pm 0.53$ ). In other words, as previously noted for WT strains, ClpX may degrade RhII in  $\Delta lasR$  cells and play a role in the  $\Delta lasR$  cell response to citrate. The distinct responses and mechanisms identified between LasR<sup>+</sup> and LasR<sup>−</sup> strains under iron limitation and exposure to the low-iron-associated molecules, citrate and pyochelin, enabled increases in antagonistic factors beyond monoculture levels as an emergent property of *P. aeruginosa* intraspecies interactions.

## DISCUSSION

In this study, we described an emergent outcome of coculturing LasR<sup>−</sup> and LasR<sup>+</sup> strains of *P. aeruginosa* in which their interactions promoted the toxic exoproducts pyocyanin and rhamnolipids (Fig. 6 provides a model). We determined that, in coculture, the  $\Delta lasR$  strain produces the siderophore pyochelin and that exogenous pyochelin induced citrate release more strongly in the WT than in  $\Delta lasR$  strain. Citrate increased RhII protein levels and induced RhIR-dependent activity only in  $\Delta lasR$  cells and not WT cells (Fig. 6). Western blot analyses of RhII-HA expressed from a regulated promoter led us to propose that the increase in RhIR-dependent signaling is due to decreased degradation of RhII by ClpXP, a known negative regulator of RhII (65, 66), or through other mechanisms of posttranscriptional regulation. The differences in siderophore production, citrate release, and RhIR/I-dependent activation between *P. aeruginosa* LasR<sup>+</sup> and LasR<sup>−</sup> strains in coculture reflect the pronounced differences between

strains that drive QS reactivation. Previous studies have shown that LasR<sup>−</sup> strain colony morphology and phenazine production change in the presence of other species such as *Candida albicans* (29) and *Staphylococcus aureus* (see Fig. 3B in reference 67), and future work will determine if pyochelin and citrate also participate in these interspecies interactions as many microbial interactions have been shown to be influenced by iron availability (68–71). Furthermore, the induction of RhIR activity that can occur in late-stationary-phase  $\Delta lasR$  cultures (30, 72) may relate to changes in iron or TCA cycle intermediates. While we found that WT-produced autoinducers, including 3OC12HSL, C4HSL, and PQS, were not required for coculture stimulation, they clearly contributed to the enhanced RhIR-dependent activity, which is consistent with intercolony QS interactions demonstrated previously (73).

The stimulatory relationship between LasR<sup>+</sup> and LasR<sup>−</sup> strains was remarkably stable as it was observed when strains were mixed within single spot colonies (Fig. 1A) and when strains were separated by either filters (Fig. 3) or millimeter distances on an agar plate (Fig. 2B). The LasR<sup>−</sup>/LasR<sup>+</sup> interactions occurred across distinct media (see Fig. S1A in the supplemental material), among genetically diverse LasR<sup>+</sup> and LasR<sup>−</sup> clinical isolates (Fig. S1B), and over a wide range of relative proportions of each type (Fig. 1C). Of note, colonies inoculated at a 80:20 WT-to- $\Delta lasR$  cell ratio had more zones with the *lasR* mutant-associated phenotypes described as sheen and lysis than colonies with a 20:80 WT-to- $\Delta lasR$  cell ratio (Fig. 1C). At both ratios,  $\Delta lasR$  cell numbers increased slightly relative to level of the wild type (Fig. 1D). We propose that the reduced appearance of sheen and lysis in mixed colonies inoculated with more  $\Delta lasR$  cells reflects a requirement for a sufficient proportion of  $\Delta lasR$  cells to initiate the WT- $\Delta lasR$  cell interactions that activate RhIR and restore a more WT-like phenotype to LasR<sup>−</sup> cells. Furthermore, if RhIR signaling is not fully activated in  $\Delta lasR$  cells, there may be regions of increased  $\Delta lasR$  cell killing via WT-produced factors such as cyanide (74).

The consequences of this intraspecies interaction may explain the worse outcomes exhibited by patients in which LasR<sup>−</sup> strains are detected (13), but future studies that include genotypes, monoculture and coculture phenotypes, and longitudinal outcome data will be required. RhIR plays other important roles in host interactions (75) which may benefit *P. aeruginosa* LasR<sup>−</sup> strains. The observation that *rhIR* mutants are rare in natural isolates and that LasR<sup>−</sup> strains with active RhIR are virulent (25, 76) underscores the relevance of this mechanism and highlights the importance of understanding how microbial interactions influence RhIR activity.

As studies of inter- and intraspecies interactions progress, it is becoming increasingly clear that the environment can dictate the outcome of microbial interactions (77). In fact, even the importance of QS regulation for fitness depends on nutrient sources and conditions (78, 79). As  $\Delta lasR$  cell-produced pyochelin was a key component of the interaction and as pyochelin production is repressed under conditions of excess iron, it was not surprising that iron supplementation suppressed the interaction (Fig. 3F and Fig. S3C and D) without significantly altering the final colony CFU count or strain ratios relative to those of the LB control (Fig. S4). Siderophore-mediated iron uptake is often required *in vivo* (34, 80, 81) due to iron sequestration by host proteins (82–85); thus, *in vivo* settings may support these interactions.

Interestingly, pyoverdine, the higher-affinity siderophore, was not required for the coculture response, mirroring findings that genes for biosynthesis of pyoverdine, but not pyochelin, are commonly disrupted in isolates from chronic CF patients (39–41). In the absence of pyoverdine (i.e., the  $\Delta lasR \Delta pvdA$  strain), we observed more pyocyanin in coculture with the WT than with the  $\Delta lasR$  strain (Fig. 3D), and we speculate that this is due to increased pyochelin production by  $\Delta pvdA$  cells, but future studies will be required to test this model. It was interesting to find that in WT/ $\Delta lasR$  coculture, heme-related proteins, *hasAP*, *hasS*, and *hasD*, were among the top eight most-upregulated genes by the  $\Delta lasR$  strain because the presence of *lasR* mutants and heme content are both reported biomarkers of disease progression in CF patients (13, 86). Coculture-induced *lasR* mutant phenotypes may link these two correlative observations.



Citrate, a TCA intermediate released under iron limitation as a result of overflow metabolism (43, 44, 60, 87), can be used by *P. aeruginosa* and other microbes for iron acquisition due to its iron chelating properties (88). The increased siderophore production by  $\Delta lasR$  cells in coculture likely reflects different metabolic strategies between genotypes. Ongoing work will investigate the mechanisms that drive differences in metabolism and iron requirements in order to determine how these differences shape microbial and host interactions. It is likely that Crc-mediated catabolite repression is involved in the response to citrate and the control of RhlI levels (64, 66). That a mechanism exists for the induction of RhlR-mediated QS in response to citrate secreted when iron is limiting dovetails with reports of increased expression of the *P. aeruginosa* QS regulon in low iron in LasR<sup>+</sup> cells (89–91). This coordinate regulation may aid in iron acquisition as QS-controlled phenazines, such as pyocyanin, reduce poorly soluble Fe<sup>3+</sup> to Fe<sup>2+</sup> and facilitate its uptake via the Feo system (92). Additionally, rhamnolipids have been employed for iron remediation (93, 94), which suggests that their surfactant activity may increase *P. aeruginosa* substrate iron uptake in part through hydroxy-alkyl-quinolone-dependent mechanisms (95).

Given that anthranilate did not alter  $\Delta lasR$  colony morphology or phenazine production, we did not further investigate anthranilate as a cross-fed metabolite involved in WT- $\Delta lasR$  cell interactions. We speculate that the increased expression of anthranilate catabolism genes in coculture may be more reflective of increases in RhlR activity than increased exposure to anthranilate given reports highlighting RhlR activation of *antABC* and *catABC* anthranilate catabolism genes (59).

As the presence of heterogeneous genotypes within single-species populations becomes increasingly appreciated, it is important to understand how commonly encountered genotypes interact to influence population-level behavior. Other work shows that cocultures can influence the survival of other genotypes (96, 97). Here, we show that intergenotype interactions lead to increased RhlR-dependent signaling in LasR<sup>−</sup> strains. It is likely that a wide array of such interactions has yet to be uncovered.

## MATERIALS AND METHODS

**Strains and growth conditions.** Strains used in this study are listed in Table S2 in the supplemental material. Bacteria were maintained on LB (lysogeny broth) medium with 1.5% agar. *Saccharomyces cerevisiae* strains for cloning were maintained on yeast-peptone-dextrose (YPD) medium with 2% agar. With the exception of pyochelin complementation experiments, which were performed in 12-well dishes with a 2-ml total volume containing 50  $\mu$ M pyochelin (or an equal-volume negative-control extract), colony biofilm assays were performed in 100-mm petri dishes with a 25-ml total volume. Where stated, a 20 mM concentration of the indicated metabolite was added to the medium (liquid or molten agar). Planktonic cultures were grown on roller drums at 37°C. Artificial sputum medium (ASM) was made as described previously (27).

**Competition assays.** Competition assays were performed to determine the relative fitness of *P. aeruginosa* mutants. Strains to be competed were grown overnight and adjusted to an optical density at 600 nm (OD<sub>600</sub>) of 1. Competing strains were combined with the PA14 *att::lacZ* strain in a 1:1 ratio unless otherwise stated. Following a 15-s vortex, 5  $\mu$ l of the combined suspension was spotted on LB agar. After 16 h, colony biofilms (and agar) were cored, placed in 1.5-ml tubes with 500  $\mu$ l of LB, and agitated vigorously for 5 min using a Genie Disruptor (Zymo). This suspension was diluted, spread on LB plates supplemented with 150  $\mu$ g/ml 5-bromo-4-chloro-3-indolyl-D-galactopyranoside (X-Gal) using glass beads, and incubated at 37°C until blue colonies were visible (~24 h). The numbers of blue and white colonies per plate were counted, and the final proportions were recorded. Each competition was run in triplicate on 3 separate days.

**Additional methods.** See Text S1 in the supplemental material for methods describing plasmid construction, pyocyanin quantification, colony proximity image analysis, swarming, RNA collection and processing, pyochelin extraction and quantification, citrate quantification,  $\beta$ -galactosidase quantification, Western blotting, and acyl-homoserine lactone activity assays.

**Data availability.** Data for RNA-seq analysis of *P. aeruginosa*  $\Delta lasR$  grown on the  $\Delta lasR$  or WT strain in coculture has been uploaded to the Gene Expression Omnibus (GEO) repository (<https://www.ncbi.nlm.nih.gov/geo/>) under accession number GSE149385.

## SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**TEXT S1**, DOCX file, 0.03 MB.

**FIG S1**, PDF file, 0.3 MB.

**FIG S2**, PDF file, 0.1 MB.

**FIG S3**, PDF file, 2.4 MB.

**FIG S4**, PDF file, 0.3 MB.

**FIG S5**, PDF file, 0.1 MB.

**FIG S6**, PDF file, 0.1 MB.

**FIG S7**, PDF file, 0.5 MB.

**TABLE S1**, XLSX file, 0.02 MB.

**TABLE S2**, DOCX file, 0.1 MB.

## ACKNOWLEDGMENTS

Research reported in this publication was supported by grants from the Cystic Fibrosis Foundation HOGAN19G0 and NIH/NIAID T32AI007519 (D.L.M.). Additional support came from NIGMS P20GM113132 through the Molecular Interactions and Imaging Core (MIIC), STANTO19R0 from the Cystic Fibrosis Foundation, and NIDDK P30-DK117469 (Dartmouth Cystic Fibrosis Research Center). RNA-seq was carried out at Dartmouth Medical School in the Genomics Shared Resource, which was established by equipment grants from the NIH and NSF and is supported in part by a Cancer Center Core Grant (P30CA023108) from the National Cancer Institute.

We also thank Georgia Doing for preliminary RNA-seq analysis, Pat Occhipinti for swapping the antibiotic marker on the *rhII*-HA expression vector, and Carla Cugini for the  $\Delta$ *lasR**clpX*::TnM mutant.

## REFERENCES

- Demers EG, Biermann AR, Masonjones S, Crocker AW, Ashare A, Stajich JE, Hogan DA. 2018. Evolution of drug resistance in an antifungal-naïve chronic *Candida lusitanae* infection. *Proc Natl Acad Sci U S A* 115: 12040–12045. <https://doi.org/10.1073/pnas.1807698115>.
- Boles BR, Thoendel M, Singh PK. 2004. Self-generated diversity produces “insurance effects” in biofilm communities. *Proc Natl Acad Sci U S A* 101:16630–16635. <https://doi.org/10.1073/pnas.0407460101>.
- Zhao S, Lieberman TD, Poyet M, Kauffman KM, Gibbons SM, Groussin M, Xavier RJ, Alm EJ. 2019. Adaptive evolution within gut microbiomes of healthy people. *Cell Host Microbe* 25:656–667.e8. <https://doi.org/10.1016/j.chom.2019.03.007>.
- Azimi S, Roberts AEL, Peng S, Weitz JS, McNally A, Brown SP, Diggle SP. 2020. Allelic polymorphism shapes community function in evolving *Pseudomonas aeruginosa* populations. *ISME J* 14:1929–1942. <https://doi.org/10.1038/s41396-020-0652-0>.
- Bos LD, Meinardi S, Blake D, Whetson K. 2016. Bacteria in the airways of patients with cystic fibrosis are genetically capable of producing VOCs in breath. *J Breath Res* 10:047103. <https://doi.org/10.1088/1752-7163/10/4/047103>.
- Jorgensen KM, Wassermann T, Johansen HK, Christiansen LE, Molin S, Hoiby N, Ciofu O. 2015. Diversity of metabolic profiles of cystic fibrosis *Pseudomonas aeruginosa* during the early stages of lung infection. *Microbiology* 161:1447–1462. <https://doi.org/10.1099/mic.0.000093>.
- Markussen T, Marvig RL, Gómez-Lozano M, Aanæs K, Burleigh AE, Høiby N, Johansen HK, Molin S, Jelsbak L. 2014. Environmental heterogeneity drives within-host diversification and evolution of *Pseudomonas aeruginosa*. *mBio* 5:e01592-14. <https://doi.org/10.1128/mBio.01592-14>.
- Wilder CN, Allada G, Schuster M. 2009. Instantaneous within-patient diversity of *Pseudomonas aeruginosa* quorum sensing populations from cystic fibrosis lung infections. *Infect Immun* 77:5631–5639. <https://doi.org/10.1128/IAI.00755-09>.
- Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* evolutionary adaptation and diversification in cystic fibrosis chronic lung infections. *Trends Microbiol* 24:327–337. <https://doi.org/10.1016/j.tim.2016.01.008>.
- Woo TE, Duong J, Jervis NM, Rabin HR, Parkins MD, Storey DG. 2016. Virulence adaptations of *Pseudomonas aeruginosa* isolated from patients with non-cystic fibrosis bronchiectasis. *Microbiology* 162:2126–2135. <https://doi.org/10.1099/mic.0.000393>.
- Workentine ML, Sibley CD, Gleason B, Purighalla S, Norgaard-Gron JC, Parkins MD, Rabin HR, Surette MG. 2013. Phenotypic heterogeneity of *Pseudomonas aeruginosa* populations in a cystic fibrosis patient. *PLoS One* 8:e60225. <https://doi.org/10.1371/journal.pone.0060225>.
- Hammond JH, Hebert WP, Naimie A, Ray K, Van Gelder RD, DiGiandomenico A, Lalitha P, Srinivasan M, Acharya NR, Lietman T, Hogan DA, Zegans ME. 2016. Environmentally endemic *Pseudomonas aeruginosa* strains with mutations in *lasR* are associated with increased disease severity in corneal ulcers. *mSphere* 1:e00140-16. <https://doi.org/10.1128/mSphere.00140-16>.
- Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW, Miller SI. 2009. *Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis lung disease progression. *J Cyst Fibros* 8:66–70. <https://doi.org/10.1016/j.jcf.2008.09.006>.
- Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. *Proc Natl Acad Sci U S A* 103: 8487–8492. <https://doi.org/10.1073/pnas.0602138103>.
- Cabrol S, Olliver A, Pier GB, Andrement A, Ruimy R. 2003. Transcription of quorum-sensing system genes in clinical and environmental isolates of *Pseudomonas aeruginosa*. *J Bacteriol* 185:7222–7230. <https://doi.org/10.1128/JB.185.24.7222-7230.2003>.
- Denervaud V, TuQuoc P, Blanc D, Favre-Bonte S, Krishnapillai V, Reimann C, Haas D, van Delden C. 2004. Characterization of cell-to-cell signaling-deficient *Pseudomonas aeruginosa* strains colonizing intubated patients. *J Clin Microbiol* 42:554–562. <https://doi.org/10.1128/jcm.42.2.554-562.2004>.
- Lee J, Zhang L. 2015. The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein Cell* 6:26–41. <https://doi.org/10.1007/s13238-014-0100-x>.
- Kanthakumar K, Taylor G, Tsang KW, Cundell DR, Rutman A, Smith S, Jeffery PK, Cole PJ, Wilson R. 1993. Mechanisms of action of *Pseudomonas aeruginosa* pyocyanin on human ciliary beat *in vitro*. *Infect Immun* 61:2848–2853. <https://doi.org/10.1128/IAI.61.7.2848-2853.1993>.
- Wilson R, Pitt T, Taylor G, Watson D, MacDermot J, Sykes D, Roberts D, Cole P. 1987. Pyocyanin and 1-hydroxyphenazine produced by *Pseudomonas aeruginosa* inhibit the beating of human respiratory cilia *in vitro*. *J Clin Invest* 79:221–229. <https://doi.org/10.1172/JCI112787>.
- Wilson R, Roberts D, Cole P. 1985. Effect of bacterial products on human ciliary function *in vitro*. *Thorax* 40:125–131. <https://doi.org/10.1136/thx.40.2.125>.
- Blumer C, Haas D. 2000. Iron regulation of the *hcnABC* genes encoding hydrogen cyanide synthase depends on the anaerobic regulator ANR rather than on the global activator GacA in *Pseudomonas fluorescens* CHAO. *Microbiology* 146:2417–2424. <https://doi.org/10.1099/00221287-146-10-2417>.
- Howe J, Bauer J, Andrä J, Schromm AB, Ernst M, Rösle M, Zähringer U, Rademann J, Brandenburg K. 2006. Biophysical characterization of syn-

- thetic rhamnolipids. *FEBS J* 273:5101–5112. <https://doi.org/10.1111/j.1742-4658.2006.05507.x>.
23. Ortiz A, Teruel JA, Espuny MJ, Marqués A, Manresa Á, Aranda FJ. 2006. Effects of dirhamnolipid on the structural properties of phosphatidylcholine membranes. *Int J Pharm* 325:99–107. <https://doi.org/10.1016/j.ijpharm.2006.06.028>.
  24. Moussa Z, Chebl M, Patra D. 2017. Interaction of curcumin with 1,2-dioctadecanoyl-sn-glycero-3-phosphocholine liposomes: intercalation of rhamnolipids enhances membrane fluidity, permeability and stability of drug molecule. *Colloids Surf B Biointerfaces* 149:30–37. <https://doi.org/10.1016/j.colsurfb.2016.10.002>.
  25. Feltner JB, Wolter DJ, Pope CE, Groleau MC, Smalley NE, Greenberg EP, Mayer-Hamblett N, Burns J, Deziel E, Hoffman LR, Dandekar AA. 2016. LasR variant cystic fibrosis isolates reveal an adaptable quorum-sensing hierarchy in *Pseudomonas aeruginosa*. *mBio* 7:e01513-16. <https://doi.org/10.1128/mBio.01513-16>.
  26. D'Argenio D, Wu M, Hoffman L, Kulasekara H, Déziel E, Smith E, Nguyen H, Ernst R, Larson Freeman T, Spencer D, Brittnacher M, Hayden H, Selgrade S, Klausen M, Goodlett D, Burns J, Ramsey B, Miller S. 2007. Growth phenotypes of *Pseudomonas aeruginosa* lasR mutants adapted to the airways of cystic fibrosis patients. *Mol Microbiol* 64:512–533. <https://doi.org/10.1111/j.1365-2958.2007.05678.x>.
  27. Clay ME, Hammond JH, Zhong F, Chen X, Kowalski CH, Lee AJ, Porter MS, Hampton TH, Greene CS, Pletneva EV, Hogan DA. 2020. *Pseudomonas aeruginosa* lasR mutant fitness in microoxia is supported by an Anr-regulated oxygen-binding hemerythrin. *Proc Natl Acad Sci U S A* 117:3167–3173. <https://doi.org/10.1073/pnas.1917576117>.
  28. Basta DW, Bergkessel M, Newman DK. 2017. Identification of fitness determinants during energy-limited growth arrest in *Pseudomonas aeruginosa*. *mBio* 8:e01170-17. <https://doi.org/10.1128/mBio.01170-17>.
  29. Cugini C, Morales DK, Hogan DA. 2010. *Candida albicans*-produced farnesol stimulates *Pseudomonas* quinolone signal production in LasR-defective *Pseudomonas aeruginosa* strains. *Microbiology* 156:3096–3107. <https://doi.org/10.1099/mic.0.037911-0>.
  30. Cabeen MT. 2014. Stationary phase-specific virulence factor overproduction by a lasR mutant of *Pseudomonas aeruginosa*. *PLoS One* 9:e88743. <https://doi.org/10.1371/journal.pone.0088743>.
  31. Van Delden C, Pesci EC, Pearson JP, Iglewski BH. 1998. Starvation selection restores elastase and rhamnolipid production in a *Pseudomonas aeruginosa* quorum-sensing mutant. *Infect Immun* 66:4499–4502. <https://doi.org/10.1128/66.9.4499-4502.1998>.
  32. Ankenbauer RG, Toyokuni T, Staley A, Rinehart KL, Cox CD. 1988. Synthesis and biological activity of pyochelin, a siderophore of *Pseudomonas aeruginosa*. *J Bacteriol* 170:5344–5351. <https://doi.org/10.1128/jb.170.11.5344-5351.1988>.
  33. Brandel J, Humbert N, Elhabiri M, Schalk IJ, Mislin GL, Albrecht-Gary AM. 2012. Pyochelin, a siderophore of *Pseudomonas aeruginosa*: physico-chemical characterization of the iron(III), copper(II) and zinc(II) complexes. *Dalton Trans* 41:2820–2834. <https://doi.org/10.1039/c1dt11804h>.
  34. Gi M, Lee KM, Kim SC, Yoon JH, Yoon SS, Choi JY. 2015. A novel siderophore system is essential for the growth of *Pseudomonas aeruginosa* in airway mucus. *Sci Rep* 5:14644. <https://doi.org/10.1038/srep14644>.
  35. Cox CD, Adams P. 1985. Siderophore activity of pyoverdine for *Pseudomonas aeruginosa*. *Infect Immun* 48:130–138. <https://doi.org/10.1128/IAI.48.1.130-138.1985>.
  36. Bhakta MN, Wilks A. 2006. The mechanism of heme transfer from the cytoplasmic heme binding protein PhuS to the  $\delta$ -regioselective heme oxygenase of *Pseudomonas aeruginosa*. *Biochemistry* 45:11642–11649. <https://doi.org/10.1021/bi060980l>.
  37. Wegele R, Tasler R, Zeng Y, Rivera M, Frankenberg-Dinkel N. 2004. The heme oxygenase(s)-phytochrome system of *Pseudomonas aeruginosa*. *J Biol Chem* 279:45791–45802. <https://doi.org/10.1074/jbc.M408303200>.
  38. Zhou H, Lu F, Latham C, Zander DS, Visner GA. 2004. Heme oxygenase-1 expression in human lungs with cystic fibrosis and cytoprotective effects against *Pseudomonas aeruginosa* *in vitro*. *Am J Respir Crit Care Med* 170:633–640. <https://doi.org/10.1164/rccm.200311-1607OC>.
  39. Nguyen AT, O'Neill MJ, Watts AM, Robson CL, Lamont IL, Wilks A, Oglesby-Sherrouse AG. 2014. Adaptation of iron homeostasis pathways by a *Pseudomonas aeruginosa* pyoverdine mutant in the cystic fibrosis lung. *J Bacteriol* 196:2265–2276. <https://doi.org/10.1128/JB.01491-14>.
  40. Marvig RL, Damkjaer S, Khademi SMH, Markussen TM, Molin S, Jelsbak L. 2014. Within-host evolution of *Pseudomonas aeruginosa* reveals adaptation toward iron acquisition from hemoglobin. *mBio* 5:e00966-14. <https://doi.org/10.1128/mBio.00966-14>.
  41. Andersen SB, Marvig RL, Molin S, Krogh Johansen H, Griffin AS. 2015. Long-term social dynamics drive loss of function in pathogenic bacteria. *Proc Natl Acad Sci U S A* 112:10756–10761. <https://doi.org/10.1073/pnas.1508324112>.
  42. Oexle H, Gnaiger E, Weiss G. 1999. Iron-dependent changes in cellular energy metabolism: influence on citric acid cycle and oxidative phosphorylation. *Biochim Biophys Acta* 1413:99–107. [https://doi.org/10.1016/S0005-2728\(99\)00088-2](https://doi.org/10.1016/S0005-2728(99)00088-2).
  43. Carlson RP, Beck AE, Phalak P, Fields MW, Gedeon T, Hanley L, Harcombe WR, Henson MA, Heys JJ. 2018. Competitive resource allocation to metabolic pathways contributes to overflow metabolisms and emergent properties in cross-feeding microbial consortia. *Biochem Soc Trans* 46:269–284. <https://doi.org/10.1042/BST20170242>.
  44. Sasnow SS, Wei H, Aristilde L. 2016. Bypasses in intracellular glucose metabolism in iron-limited *Pseudomonas putida*. *Microbiologyopen* 5:3–20. <https://doi.org/10.1002/mbo3.287>.
  45. Hammond JH, Dolben EF, Smith TJ, Bhujji S, Hogan DA. 2015. Links between Anr and quorum sensing in *Pseudomonas aeruginosa* biofilms. *J Bacteriol* 197:2810–2820. <https://doi.org/10.1128/JB.00182-15>.
  46. O'Loughlin CT, Miller LC, Siryaporn A, Drescher K, Semmelhack MF, Bassler BL. 2013. A quorum-sensing inhibitor blocks *Pseudomonas aeruginosa* virulence and biofilm formation. *Proc Natl Acad Sci U S A* 110:17981–17986. <https://doi.org/10.1073/pnas.1316981110>.
  47. Lau GW, Hassett DJ, Ran H, Kong F. 2004. The role of pyocyanin in *Pseudomonas aeruginosa* infection. *Trends Mol Med* 10:599–606. <https://doi.org/10.1016/j.molmed.2004.10.002>.
  48. Whiteley M, Lee KM, Greenberg EP. 1999. Identification of genes controlled by quorum sensing in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 96:13904–13909. <https://doi.org/10.1073/pnas.96.24.13904>.
  49. Brint JM, Ohman DE. 1995. Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RhlR-RhlI, another set of regulators in strain PAO1 with homology to the autoinducer-responsive LuxR-LuxI family. *J Bacteriol* 177:7155–7163. <https://doi.org/10.1128/jb.177.24.7155-7163.1995>.
  50. Rahme LG, Tan MW, Le L, Wong SM, Tompkins RG, Calderwood SB, Ausubel FM. 1997. Use of model plant hosts to identify *Pseudomonas aeruginosa* virulence factors. *Proc Natl Acad Sci U S A* 94:13245–13250. <https://doi.org/10.1073/pnas.94.24.13245>.
  51. Köhler T, Curty LK, Barja F, van Delden C, Pechère JC. 2000. Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. *J Bacteriol* 182:5990–5996. <https://doi.org/10.1128/jb.182.21.5990-5996.2000>.
  52. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. 2019. PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Res* 47:D419–D426. <https://doi.org/10.1093/nar/gky1038>.
  53. Loudon BC, Haarmann D, Lynne AM. 2011. Use of blue agar CAS assay for siderophore detection. *J Microbiol Biol Educ* 12:51–53. <https://doi.org/10.1128/jmbe.v12i1.249>.
  54. Calfee MW, Coleman JP, Pesci EC. 2001. Interference with *Pseudomonas* quinolone signal synthesis inhibits virulence factor expression by *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 98:11633–11637. <https://doi.org/10.1073/pnas.201328498>.
  55. Farrow JM, 3rd, Pesci EC. 2007. Two distinct pathways supply anthranilate as a precursor of the *Pseudomonas* quinolone signal. *J Bacteriol* 189:3425–3433. <https://doi.org/10.1128/JB.00209-07>.
  56. The UniProt Consortium. 2018. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 47:D506–D515. <https://doi.org/10.1093/nar/gky1049>.
  57. Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. 2016. Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. *Nucleic Acids Res* 44:D646–D653. <https://doi.org/10.1093/nar/gkv1227>.
  58. Lesuisse E, Simon M, Klein R, Labbe P. 1992. Excretion of anthranilate and 3-hydroxyanthranilate by *Saccharomyces cerevisiae*: relationship to iron metabolism. *J Gen Microbiol* 138:85–89. <https://doi.org/10.1099/00221287-138-1-85>.
  59. Choi Y, Park HY, Park SJ, Park SJ, Kim SK, Ha C, Im SJ, Lee JH. 2011. Growth phase-differential quorum sensing regulation of anthranilate metabolism in *Pseudomonas aeruginosa*. *Mol Cells* 32:57–65. <https://doi.org/10.1007/s10059-011-2322-6>.
  60. Odoni DI, van Gaal MP, Schonewille T, Tamayo-Ramos JA, Martins dos



- Santos VAP, Suarez-Diez M, Schaap PJ. 2017. *Aspergillus niger* secretes citrate to increase iron bioavailability. *Front Microbiol* 8:1424. <https://doi.org/10.3389/fmicb.2017.01424>.
61. Valentini M, Storelli N, Lapouge K. 2011. Identification of C(4)-dicarboxylate transport systems in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* 193:4307–4316. <https://doi.org/10.1128/JB.05074-11>.
  62. Filiatrault MJ, Tomblin G, Wagner VE, Van Alst N, Rumbaugh K, Sokol P, Schwengel J, Iglewski BH. 2013. *Pseudomonas aeruginosa* PA1006, which plays a role in molybdenum homeostasis, is required for nitrate utilization, biofilm formation, and virulence. *PLoS One* 8:e55594. <https://doi.org/10.1371/journal.pone.0055594>.
  63. Thomason MK, Voichek M, Dar D, Addis V, Fitzgerald D, Gottesman S, Sorek R, Greenberg EP. 2019. A *rhlI* 5' UTR-derived sRNA regulates RhlR-dependent quorum sensing in *Pseudomonas aeruginosa*. *mBio* 10:e02253-19. <https://doi.org/10.1128/mBio.02253-19>.
  64. Yang N, Ding S, Chen F, Zhang X, Xia Y, Di H, Cao Q, Deng X, Wu M, Wong CC, Tian XX, Yang CG, Zhao J, Lan L. 2015. The Crc protein participates in down-regulation of the *Lon* gene to promote rhamnolipid production and Rhl quorum sensing in *Pseudomonas aeruginosa*. *Mol Microbiol* 96:526–547. <https://doi.org/10.1111/mmi.12954>.
  65. Takaya A, Tabuchi F, Tsuchiya H, Isogai E, Yamamoto T. 2008. Negative regulation of quorum-sensing systems in *Pseudomonas aeruginosa* by ATP-dependent *Lon* protease. *J Bacteriol* 190:4181–4188. <https://doi.org/10.1128/JB.01873-07>.
  66. Yang N, Lan L. 2016. *Pseudomonas aeruginosa* *Lon* and *ClpXP* proteases: roles in linking carbon catabolite repression system with quorum-sensing system. *Curr Genet* 62:1–6. <https://doi.org/10.1007/s00294-015-0499-5>.
  67. Hoffman LR, Richardson AR, Houston LS, Kulasekara HD, Martens-Habbena W, Klausen M, Burns JL, Stahl DA, Hassett DJ, Fang FC, Miller SI. 2010. Nutrient availability as a mechanism for selection of antibiotic tolerant *Pseudomonas aeruginosa* within the CF airway. *PLoS Pathog* 6:e1000712. <https://doi.org/10.1371/journal.ppat.1000712>.
  68. Harrison F, Paul J, Massey RC, Buckling A. 2008. Interspecific competition and siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *ISME J* 2:49–55. <https://doi.org/10.1038/ismej.2007.96>.
  69. Lopez-Medina E, Fan D, Coughlin LA, Ho EX, Lamont IL, Reimann C, Hooper LV, Koh AY. 2015. *Candida albicans* inhibits *Pseudomonas aeruginosa* virulence through suppression of pyochelin and pyoverdine biosynthesis. *PLoS Pathog* 11:e1005129. <https://doi.org/10.1371/journal.ppat.1005129>.
  70. Weaver VB, Kolter R. 2004. *Burkholderia* spp. alter *Pseudomonas aeruginosa* physiology through iron sequestration. *J Bacteriol* 186:2376–2384. <https://doi.org/10.1128/jb.186.8.2376-2384.2004>.
  71. Scott JE, Li K, Filkins LM, Zhu B, Kuchma SL, Schwartzman JD, O'Toole GA. 2019. *Pseudomonas aeruginosa* can inhibit growth of *Streptococcal* species via siderophore production. *J Bacteriol* 201:e00014-19. <https://doi.org/10.1128/JB.00014-19>.
  72. Dekimpe V, Deziel E. 2009. Revisiting the quorum-sensing hierarchy in *Pseudomonas aeruginosa*: the transcriptional regulator RhlR regulates LasR-specific factors. *Microbiology* 155:712–723. <https://doi.org/10.1099/mic.0.022764-0>.
  73. Darch SE, Simoska O, Fitzpatrick M, Barraza JP, Stevenson KJ, Bonnezaze RT, Shear JB, Whiteley M. 2018. Spatial determinants of quorum signaling in a *Pseudomonas aeruginosa* infection model. *Proc Natl Acad Sci U S A* 115:4779–4784. <https://doi.org/10.1073/pnas.1719317115>.
  74. Wang M, Schaefer AL, Dandekar AA, Greenberg EP. 2015. Quorum sensing and policing of *Pseudomonas aeruginosa* social cheaters. *Proc Natl Acad Sci U S A* 112:2187–2191. <https://doi.org/10.1073/pnas.1500704112>.
  75. Haller S, Franchet A, Hakkim A, Chen J, Drenkard E, Yu S, Schirmeier S, Li Z, Martins N, Ausubel FM, Liegeois S, Ferrandon D. 2018. Quorum-sensing regulator RhlR but not its autoinducer RhlI enables *Pseudomonas* to evade opsonization. *EMBO Rep* 19:e44880. <https://doi.org/10.15252/embr.201744880>.
  76. Cruz RL, Asfahl KL, Van den Bossche S, Coenye T, Crabbé A, Dandekar AA. 2020. RhlR-regulated acyl-homoserine lactone quorum sensing in a cystic fibrosis isolate of *Pseudomonas aeruginosa*. *mBio* 11:e00532-20. <https://doi.org/10.1128/mBio.00532-20>.
  77. Doing G, Koeppen J, Occipinti P, Hogan D. 2020. Conditional antagonism in co-cultures of *Pseudomonas aeruginosa* and *Candida albicans*: an intersection of ethanol and phosphate signaling distilled from dual-seq transcriptomics. *bioRxiv* <https://www.biorxiv.org/content/10.1101/2020.04.20.050765v2>.
  78. Dandekar AA, Chugani S, Greenberg EP. 2012. Bacterial quorum sensing and metabolic incentives to cooperate. *Science* 338:264–266. <https://doi.org/10.1126/science.1227289>.
  79. Mund A, Diggle SP, Harrison F. 2017. The fitness of *Pseudomonas aeruginosa* quorum sensing signal cheats is influenced by the diffusivity of the environment. *mBio* 8:e00816-17. <https://doi.org/10.1128/mBio.00816-17>.
  80. Reid DW, Anderson GJ, Lamont IL. 2009. Role of lung iron in determining the bacterial and host struggle in cystic fibrosis. *Am J Physiol Lung Cell Mol Physiol* 297:L795–L802. <https://doi.org/10.1152/ajplung.00132.2009>.
  81. Lamont IL, Konings AF, Reid DW. 2009. Iron acquisition by *Pseudomonas aeruginosa* in the lungs of patients with cystic fibrosis. *Biometals* 22:53–60. <https://doi.org/10.1007/s10534-008-9197-9>.
  82. Britigan BE, Rasmussen GT, Olakanmi O, Cox CD. 2000. Iron acquisition from *Pseudomonas aeruginosa* siderophores by human phagocytes: an additional mechanism of host defense through iron sequestration? *Infect Immun* 68:1271–1275. <https://doi.org/10.1128/iai.68.3.1271-1275.2000>.
  83. Nakashige TG, Zhang B, Krebs C, Nolan EM. 2015. Human calprotectin is an iron-sequestering host-defense protein. *Nat Chem Biol* 11:765–771. <https://doi.org/10.1038/nchembio.1891>.
  84. Ward PP, Conneely OM. 2004. Lactoferrin: role in iron homeostasis and host defense against microbial infection. *Biometals* 17:203–208. <https://doi.org/10.1023/b:biom.0000027693.60932.26>.
  85. Michels KR, Zhang Z, Bettina AM, Cagnina RE, Stefanova D, Burdick MD, Vaulton S, Nemeth E, Ganz T, Mehrad B. 2017. Hepcidin-mediated iron sequestration protects against bacterial dissemination during pneumonia. *JCI Insight* 2:e92002. <https://doi.org/10.1172/jci.insight.92002>.
  86. Glasser NR, Hunter RC, Liou TG, Newman DK, Mountain West CF Consortium Investigators. 2019. Refinement of metabolite detection in cystic fibrosis sputum reveals heme correlates with lung function decline. *PLoS One* 14:e0226578. <https://doi.org/10.1371/journal.pone.0226578>.
  87. Folsom JP, Parker AE, Carlson RP. 2014. Physiological and proteomic analysis of *Escherichia coli* iron-limited chemostat growth. *J Bacteriol* 196:2748–2761. <https://doi.org/10.1128/JB.01606-14>.
  88. Marshall B, Stintzi A, Gilmour C, Meyer JM, Poole K. 2009. Citrate-mediated iron uptake in *Pseudomonas aeruginosa*: involvement of the citrate-inducible FecA receptor and the FeoB ferrous iron transporter. *Microbiology* 155:305–315. <https://doi.org/10.1099/mic.0.023531-0>.
  89. Oglesby AG, Farrow JM, 3rd, Lee JH, Tomaras AP, Greenberg EP, Pesci EC, Vasil ML. 2008. The influence of iron on *Pseudomonas aeruginosa* physiology: a regulatory link between iron and quorum sensing. *J Biol Chem* 283:15558–15567. <https://doi.org/10.1074/jbc.M707840200>.
  90. Kim EJ, Sabra W, Zeng AP. 2003. Iron deficiency leads to inhibition of oxygen transfer and enhanced formation of virulence factors in cultures of *Pseudomonas aeruginosa* PAO1. *Microbiology* 149:2627–2634. <https://doi.org/10.1099/mic.0.26276-0>.
  91. Kim EJ, Wang W, Deckwer WD, Zeng AP. 2005. Expression of the quorum-sensing regulatory protein LasR is strongly affected by iron and oxygen concentrations in cultures of *Pseudomonas aeruginosa* irrespective of cell density. *Microbiology* 151:1127–1138. <https://doi.org/10.1099/mic.0.27566-0>.
  92. Cox CD. 1986. Role of pyocyanin in the acquisition of iron from transferrin. *Infect Immun* 52:263–270. <https://doi.org/10.1128/IAI.52.1.263-270.1986>.
  93. Akintunde TA, Abioye P, Oyeleke SB, Boboye BE, Ijah UJ. 2015. Remediation of iron using rhamnolipid-surfactant produced by *Pseudomonas aeruginosa*. *Res J Environ Sci* 9:169–177. <https://doi.org/10.3923/rjes.2015.169.177>.
  94. Wang S, Mulligan CN. 2009. Rhamnolipid biosurfactant-enhanced soil flushing for the removal of arsenic and heavy metals from mine tailings. *Process Biochem* 44:296–301. <https://doi.org/10.1016/j.procbio.2008.11.006>.
  95. Diggle SP, Matthijs S, Wright VJ, Fletcher MP, Chhabra SR, Lamont IL, Kong X, Hider RC, Cornelis P, Camara M, Williams P. 2007. The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem Biol* 14:87–96. <https://doi.org/10.1016/j.chembiol.2006.11.014>.
  96. Malhotra S, Limoli DH, English AE, Parsek MR, Wozniak DJ. 2018. Mixed communities of mucoid and nonmucoid *Pseudomonas aeruginosa* exhibit enhanced resistance to host antimicrobials. *mBio* 9:e00275-18. <https://doi.org/10.1128/mBio.00275-18>.
  97. Oluyombo O, Penfold CN, Diggle SP. 2019. Competition in biofilms between Cystic Fibrosis isolates of *Pseudomonas aeruginosa* is shaped by R-Pyocins. *mBio* 10:e01828-18. <https://doi.org/10.1128/mBio.01828-18>.